

CLASSIFICATION OF QUADRUPLE GALOIS CANONICAL COVERS I

FRANCISCO JAVIER GALLEGO
AND
BANGERE P. PURNAPRAJNA

ABSTRACT. In this article we classify quadruple Galois canonical covers of smooth surfaces of minimal degree. The classification shows that they are either non-simple cyclic covers or bi-double covers. If they are bi-double then they are all fiber products of double covers. We construct examples to show that all the possibilities in the classification do exist. There are implications of this classification that include the existence of families with unbounded geometric genus and families with unbounded irregularity, in sharp contrast with the case of double and triple canonical covers. Together with the results of Horikawa and Konno for double and triple covers, a pattern emerges that motivates some general questions on the existence of higher degree canonical covers, some of which are answered in this article.

DEDICATED TO IGNACIO SOLS.

INTRODUCTION

Classification problems are of central importance in Algebraic Geometry. In the realm of algebraic surfaces, the geography of surfaces of general type, by far the largest class of surfaces, is much less charted and understood. An important sub-class of surfaces of general type are those whose canonical map is a cover of a simpler surface, most notably, a cover of a surface of minimal degree. In the seventies and eighties Horikawa and Konno (see [Ho1], [Ho3] and [K]) classified these covers when the degree of the cover is 2 and 3. In this article and in its sequel [GP4] we classify surfaces of general type whose canonical map is a quadruple Galois cover of a surface of minimal degree.

Covers of varieties of minimal degree have a ubiquitous presence in various contexts. They appear in the classification of surfaces of general type X with small K_X^2 done by Horikawa (see [Ho1], [Ho2], [Ho3] and [Ho4]) and play an important role in mapping the

2000 *Mathematics Subject Classification*: 14J10, 14J26, 14J29

The first author was partially supported by MCT project number BFM2000-0621. He is grateful for the hospitality of the Department of Mathematics of the University of Kansas at Lawrence. The second author is grateful to NSA for supporting this research project. He is also grateful for the hospitality of the Departamento de Álgebra of the Universidad Complutense de Madrid

geography of surfaces of general type. They are also the chief source in constructing new examples of surfaces of general type as the work of various geometers illustrates (see also [K], [MP] and [T]). These covers occur as well in the study of linear series on important threefolds such as Calabi-Yau threefolds as the work in [BS], [GP1] and [OP] shows. They also become relevant in the study of the canonical ring of a variety of general type as can be seen in results from [GP2] and [Gr] (see [Pu] for further motivation).

Compared to the canonical morphism of a curve, the canonical morphism of a surface is much more subtle and allows a much wider range of possibilities due to the existence of higher degree covers. The degree of the canonical morphism of a curve is bounded by 2. In contrast, Beauville proved that the degree of the canonical morphism from a surface of general type X onto a surface of minimal degree, or more generally, onto a surface with geometric genus $p_g = 0$, is bounded by 9 if $\chi(X) \geq 31$ (see [Be]; see also [Ga], where the bound is reduced to 8 if $p_g(X) \geq 132$).

Surfaces of minimal degree are classically known to be linear $\mathbf{P}^2$, the Veronese surface in $\mathbf{P}^5$ and rational normal scrolls, which can be smooth (these include the smooth quadric hypersurface in $\mathbf{P}^3$) or singular (these are cones over a rational normal curve). As pointed out before the classification of the canonical covers of these surfaces is only complete when the degree of the cover is 2 (see [Ho1]) and 3 (see [Ho3] and [K]). Horikawa also studied quadruple covers of linear $\mathbf{P}^2$. The next step in this classification is the study of quadruple covers of an arbitrary surface of minimal degree.

In this work we classify all quadruple Galois canonical covers of smooth surfaces of minimal degree W . In [GP4] we classify quadruple Galois covers of W when W is singular. There are many interesting consequences of the classification done here and in [GP4]. Our classification yields, among other things, some striking contrasts with double and triple covers. Before we look at them we state the main result of this article:

Theorem. *Let X be a canonical surface and let W be a smooth surface of minimal degree. If the canonical bundle of X is base-point-free and $X \xrightarrow{\varphi} W$ is a quadruple Galois canonical cover, then W is either linear $\mathbf{P}^2$ or a smooth rational normal scroll $S(m - e, m)$ with $0 \leq e \leq 2$ and $m \geq e + 1$. Let G be the Galois group of φ .*

A) *If $G = \mathbf{Z}_2 \times \mathbf{Z}_2$, then X is the fiber product over W of two double covers of W branched along divisors D_1 and D_2 and φ is the natural morphism from the fiber product to W . The divisors D_1 and D_2 satisfy:*

1) *If W is linear $\mathbf{P}^2$, then D_1 and D_2 are quartic curves.*

2) *If $W = S(m - e, m)$, then either*

2.1) *$D_1 \sim 2C_0 + (2m + 2)f$ and $D_2 \sim 4C_0 + (2e + 2)f$; or*

$e = 0$ and

2.2) *$D_1 \sim (2m + 2)f$ and $D_2 \sim 6C_0 + 2f$; or*

2.3) *$D_1 \sim (2m + 4)f$ and $D_2 \sim 6C_0$; or*

2.4) *$D_1 \sim 2C_0 + (2m + 4)f$ and $D_2 \sim 4C_0$.*

B) If $G = \mathbf{Z}_4$, then φ is the composition of two double covers $X_1 \xrightarrow{p_1} Y$ branched along a divisor D_2 and $X \xrightarrow{p_2} X_1$, branched along the ramification of p_1 and $p_1^*D_1$ and with trace zero module $p_1^*\mathcal{O}_W(-\frac{1}{2}D_1 - \frac{1}{4}D_2)$. The divisors D_1 and D_2 satisfy:

1) If W is linear $\mathbf{P}^2$, D_1 is a conic and D_2 is a quartic curve.

2) If $W = S(m - e, m)$, then either

2.1) $D_1 \sim (2m - e + 1)f$ and $D_2 \sim 4C_0 + (2e + 2)f$; or

$e = 0$, and

2.2) $D_1 \sim (2m + 4)f$, $D_2 \sim 4C_0$; or

2.3) $D_1 \sim 3C_0$, $D_2 \sim 2C_0 + 4f$ and $m = 1$.

Conversely, if $X \xrightarrow{\varphi} W$ is either

I. the fiber product over W of two double covers $X_1 \xrightarrow{p_1} W$ and $X_2 \xrightarrow{p_2} W$, branched respectively along divisors D_2 and D_1 as described in A.1, A.2.1, A.2.2, A.2.3 or A.2.4; or

II. the composition of two double covers $X_1 \xrightarrow{p_1} Y$ branched along a divisor D_2 and $X \xrightarrow{p_2} X_1$, branched along the ramification of p_1 and $p_1^*D_1$ and with trace zero module $p_1^*\mathcal{O}_W(-\frac{1}{2}D_1 - \frac{1}{4}D_2)$, with D_1 and D_2 as described in B.1, B.2.1, B.2.2 or B.2.3,

then $X \xrightarrow{\varphi} W$ is a Galois canonical cover whose Galois group is $\mathbf{Z}_2 \times \mathbf{Z}_2$ in case I and $\mathbf{Z}_4$ in case II.

The above result is split inside the paper into Corollary 1.7 and Theorems 2.11, 3.5, 3.6, 3.8 and 3.9. We also construct families of examples to show the existence of all the cases that appear in the classification. A notable fact that is not mentioned in the statement of the theorem is that there exist families of smooth surfaces of general type X for every case A.1, A.2.1, A.2.2, A.2.3 and A.2.4, (that is, all cases where $G = \mathbf{Z}_2 \times \mathbf{Z}_2$). In comparison, we show that the quadruple cyclic canonical covers of smooth surfaces of minimal degree are always singular. We also show that quadruple cyclic canonical covers are non-simple cyclic.

One of the important implications of the main result of this article is the existence of families of quadruple canonical covers with unbounded geometric genus and the existence of families with unbounded irregularity. The unboundedness of the geometric genus is in sharp contrast with the situation of triple covers and the unboundedness of the irregularity is in sharp contrast with the situation of both double and triple covers. The geometric genus of canonical double covers is unbounded but they are all regular, and even simply connected surfaces. The geometric genus of canonical triple covers is bounded by 5 (see [K]) and are all regular. For quadruple Galois covers we show the existence of families of surfaces X for each possible value of $p_g(X)$. Regarding irregularity, the classification shows that most of the quadruple Galois covers are regular. However, we do show that there exist families of surfaces with irregularity 1 (cases A.2.4, B.2.2 and B.2.3) and, most importantly, families of surfaces X for each possible value of $q(X)$ (cases A.2.2 and A.2.3).

The classification of quadruple covers provides other significant contrast with double and triple covers and brings out clearly the marked difference between even and odd degree

covers. The only smooth targets of quadruple Galois canonical covers that occur are linear $\mathbf{P}^2$ and rational normal scrolls which correspond to only three Hirzebruch surfaces, namely $\mathbf{F}_0$, $\mathbf{F}_1$ and $\mathbf{F}_2$. In the case of canonical double covers, linear $\mathbf{P}^2$ and smooth rational scrolls corresponding to every Hirzebruch surface appear as image of the canonical morphism. In the case of canonical triple covers, the list is reduced drastically and the only possible smooth target is linear $\mathbf{P}^2$ (see also [GP2], Proposition 3.3).

The classification of quadruple covers of singular targets (i.e., a cone over a rational normal curve) in [GP4] together with the results of Horikawa and Konno exhibit a striking pattern. Indeed we show that these form a bounded family with respect to both the geometric genus and irregularity as in the case of double and triples covers of singular targets. The results in this article and [GP4] predict a precise numerology that might hold for higher degree covers. The following facts make it clear what we mean: there do not exist canonical covers of odd degree of smooth scrolls (see [GP2], Proposition 3.3) and there do not exist Galois canonical covers $X \xrightarrow{\varphi} W$ of prime degree $p > 3$ of surfaces W of minimal degree if X is regular or if W is smooth (see Theorem 3.2, Corollary 3.3 and [GP2], Corollary 3.2). This motivates us to pose a general question (see Question 3.4) on the non-existence of higher, prime degree Galois canonical covers of surfaces of minimal degree.

The classification obtained in this article and in [GP4] has further applications. In [GP5], we determine the ring generators of the quadruple covers classified here and in [GP4]. The results of [GP5] show that quadruple covers serve as examples and counter examples to some questions on graded rings and normal generation of linear systems on an algebraic surface.

Acknowledgements: We thank N. Mohan Kumar for generously sharing his time for some useful discussions and insightful comments. We are very grateful to Rita Pardini, who, after seeing an earlier version of this work, remarked that we had not treated the case of non-simple cyclic covers. It was after her comment that we proved the theorems on cyclic covers. Finally, we thank the referee of the announcement [GP3] of these results, who kindly made many suggestions, not only for the announcement, but also to improve the exposition of the present article and [GP4].

1. QUADRUPLE CANONICAL COVERS OF SURFACES OF MINIMAL DEGREE

Convention: We work over an algebraically closed field of characteristic 0.

Notation: *We will follow these conventions:*

- 1) Throughout this article, unless otherwise stated, W will be an embedded smooth projective algebraic surface of minimal degree, i.e., whose degree is equal to its codimension in projective space plus 1.
- 2) Throughout this article, unless otherwise stated, X will be a projective algebraic normal surface with at worst canonical singularities. We will denote by ω_X the canonical bundle of X .

We recall the following standard notation:

3) By $\mathbf{F}_e$ we denote the Hirzebruch surface whose minimal section have self-intersection $-e$. If $e > 0$ let C_0 denote the minimal section of $\mathbf{F}_e$ and let f be one of the fibers of $\mathbf{F}_e$. If $e = 0$, C_0 will be a fiber of one of the families of lines and f will be a fiber of the other family of lines.

4) If a, b are integers such that $0 < a \leq b$, consider two disjoint linear subspaces $\mathbf{P}^a$ and $\mathbf{P}^b$ of $\mathbf{P}^{a+b+1}$. We denote by $S(a, b)$ the smooth rational normal scroll obtained by joining corresponding points of a rational normal curve in $\mathbf{P}^a$ and a rational normal curve of $\mathbf{P}^b$. Recall that $S(a, b)$ is the image of $\mathbf{F}_e$ by the embedding induced by the complete linear series $|C_0 + mf|$, with $a = m - e$, $b = m$ and $m \geq e + 1$. If $a = b$, the linear series $|mC_0 + f|$ also gives a minimal degree embedding of $\mathbf{F}_0$, equivalent to the previous one by the automorphism of $\mathbf{P}^1 \times \mathbf{P}^1 = \mathbf{F}_0$ swapping the factors. In this case *our convention will always be to choose C_0 and f so that, when W is a smooth rational normal scroll, and W is embedded by $|C_0 + mf|$.*

In this section we prove general results regarding quadruple canonical covers of surfaces of minimal degree:

Definition 1.1. *Let X and W be as in the previous notations. We will say that a surjective morphism $X \xrightarrow{\varphi} W$ is a canonical cover of W if X is surface of general type whose canonical bundle ω_X is ample and base-point-free and φ is the canonical morphism of X .*

Remark 1.2. *Although we have assumed X to have canonical singularities, some results hold in greater generality. Precisely, if for the purpose of this remark we ignore Notation 2) above and X is assumed to be a normal, locally Gorenstein surface instead then Definition 1.1 still makes sense and Theorems 2.11, 3.5, 3.6, 3.8 and 3.9 hold. We can further relax the hypotheses on X in the converse parts of Theorems 2.11, 3.5, 3.6, 3.8 and 3.9 and they hold if X is just assumed to be smooth in codimension 1, since in that case these covers are Gorenstein.*

Proposition 1.3. *Let $X \xrightarrow{\varphi} W$ be a quadruple canonical cover of W . Let $H = \mathcal{O}_W(1)$. Then $\varphi_*\mathcal{O}_X$ is a vector bundle on W and*

$$\varphi_*\mathcal{O}_X = \mathcal{O}_W \oplus E \oplus (\omega_W \otimes H^*)$$

with E vector bundle over W of rank 2. If in addition $\varphi_\mathcal{O}_X$ splits as sum of line bundles, then*

$$\varphi_*\mathcal{O}_X = \mathcal{O}_W \oplus L_1^* \oplus L_2^* \oplus (\omega_W \otimes H^*)$$

with $L_1^ \otimes L_2^* = \omega_W \otimes H^*$.*

Proof. Recall that by Definition 1.1 φ is finite, W is smooth and X is locally Cohen-Macaulay. Then φ is flat and hence $\varphi_*\mathcal{O}_X$ is a vector bundle over $\mathcal{O}_W$ of rank 4. Moreover (see [HM]), $\varphi_*\mathcal{O}_X = \mathcal{O}_W \oplus E'$, where E' is the trace zero module of φ . From relative duality we have

$$\varphi_*\omega_X = (\varphi_*\mathcal{O}_X)^* \otimes \omega_W .$$

On the other hand, by hypothesis, $\omega_X = \varphi^*H$, hence, by projection formula,

$$\varphi_*\omega_X = \varphi_*\mathcal{O}_X \otimes H .$$

Then

$$\omega_W \oplus (\omega_W \otimes (E')^*) = H \oplus (E' \otimes H) .$$

Since $\omega_W = H$ is not possible, for W is a rational surface, $E' = E \oplus (\omega_W \otimes H^*)$, with E vector bundle of rank 2. If $\varphi_*\mathcal{O}_X$ splits, let $E = L_1^* \oplus L_2^*$. Then

$$\begin{aligned} \omega_W \oplus (\omega_W \otimes L_1) \oplus (\omega_W \otimes L_2) \oplus H = \\ H \oplus (H \otimes L_1^*) \oplus (H \otimes L_2^*) \oplus \omega_W . \end{aligned}$$

Then taking the determinant of both sides of the equality gives $(L_1^* \otimes L_2^*)^{\otimes 2} = (\omega_W \otimes H^*)^{\otimes 2}$. Since W is either $\mathbf{P}^2$ or a Hirzebruch surface, then $L_1^* \otimes L_2^* = \omega_W \otimes H^*$. $\square$

Now we study in more detail the possible splittings of $\varphi_*\mathcal{O}_X$ depending on what surface W is. We start with this observation about linear $\mathbf{P}^2$:

Proposition 1.4. *Let $X \xrightarrow{\varphi} W$ be a canonical cover. If W is linear $\mathbf{P}^2$, then X is regular if and only if $\varphi_*\mathcal{O}_W$ splits as direct sum of line bundles.*

Proof. By Proposition 1.3 we know that

$$\varphi_*\mathcal{O}_X = \mathcal{O}_W \oplus E \oplus \omega_W(-1) .$$

Since $W = \mathbf{P}^2$, the intermediate cohomology of line bundles on W vanishes so by projection formula $H^1(\varphi^*\mathcal{O}_W(k)) = H^1(E(k))$. By Kodaira vanishing and duality $H^1(\varphi^*\mathcal{O}_W(k)) = 0$ except maybe if $k = 0, 1$. Then X is regular if and only if $H^1(E(k)) = 0$ for all k , and by Horrocks's splitting Criterion, this is equivalent to the splitting of E . Thus X is regular if and only if $\varphi_*\mathcal{O}_X$ splits as direct sum of line bundles. $\square$

The following proposition tells how the restriction of $\varphi_*\mathcal{O}_X$ to a smooth curve in $|\omega_X|$ splits:

Proposition 1.5. *Let W be a surface of minimal degree r , not necessarily smooth, let $X \xrightarrow{\varphi} W$ be a canonical cover of degree 4 and let C be a general smooth irreducible curve in $|\mathcal{O}_W(1)|$. If X is regular, then*

$$(\varphi_*\mathcal{O}_X)|_C = \mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-r-1) \oplus \mathcal{O}_{\mathbf{P}^1}(-r-1) \oplus \mathcal{O}_{\mathbf{P}^1}(-2r-2) .$$

Proof. See [GP2], Lemma 2.3 for details. $\square$

Finally we describe more accurately the splitting of $\varphi_*\mathcal{O}_X$ in the case X regular:

Proposition 1.6. *Let $X \xrightarrow{\varphi} W$ be a quadruple canonical cover. If $\varphi_*\mathcal{O}_X$ splits as a direct sum of line bundles, then:*

1) *If W is linear $\mathbf{P}^2$, then*

$$\varphi_*\mathcal{O}_X = \mathcal{O}_{\mathbf{P}^2} \oplus \mathcal{O}_{\mathbf{P}^2}(-2) \oplus \mathcal{O}_{\mathbf{P}^2}(-2) \oplus \mathcal{O}_{\mathbf{P}^2}(-4) ;$$

2) *W is not the Veronese surface; and*

3) *if W is a rational normal scroll and X is regular, then*

$$\begin{aligned} \varphi_*\mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-C_0 - (m+1)f) \oplus \\ &\mathcal{O}_W(-2C_0 - (e+1)f) \oplus \mathcal{O}_W(-3C_0 - (m+e+2)f) , \end{aligned}$$

where $2m - e$ is the degree of W .

Proof. To prove 1) recall that by Proposition 1.4 X is regular. Then by Proposition 1.5 the restriction of $\varphi_*\mathcal{O}_X$ to a line of W is

$$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-2) \oplus \mathcal{O}_{\mathbf{P}^1}(-2) \oplus \mathcal{O}_{\mathbf{P}^1}(-4)$$

so 1) is clear.

For 2) W is isomorphic to $\mathbf{P}^2$ and $\mathcal{O}_W(1) = \mathcal{O}_{\mathbf{P}^2}(2)$. Then by Proposition 1.3 $\varphi_*\mathcal{O}_X = \mathcal{O}_{\mathbf{P}^2} \oplus E \oplus \mathcal{O}_{\mathbf{P}^2}(-5)$. Let C be a smooth conic in W . If X is irregular, then by Proposition 1.4, $\varphi_*\mathcal{O}_X$ cannot split completely. If X is regular, then according to Proposition 1.5

$$(\varphi_*\mathcal{O}_X)_C = \mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-5) \oplus \mathcal{O}_{\mathbf{P}^1}(-5) \oplus \mathcal{O}_{\mathbf{P}^1}(-10) .$$

Then if $E = L_1^* \oplus L_2^*$ with L_1 and L_2 line bundles, $L_1^*|_C = L_2^*|_C = \mathcal{O}_{\mathbf{P}^1}(-5)$, but since C is a conic the degrees of $L_1|_C$ and $L_2|_C$ are even integers, so we get a contradiction.

For 3) recall that $\omega_W = \mathcal{O}_W(-2C_0 - (e+2)f)$ and that $\mathcal{O}_W(1) = \mathcal{O}_W(C_0 + mf)$. Then it follows by assumption and Proposition 1.3 that

$$\varphi_*\mathcal{O}_X = \mathcal{O}_W \oplus L_1^* \oplus L_2^* \oplus \mathcal{O}_W(-3C_0 - (m+e+2)f) ,$$

and $L_1 \otimes L_2 = \mathcal{O}_W(3C_0 + (m+e+2)f)$. Then, if we set $L_1 = \mathcal{O}_W(a_1C_0 + b_1f)$ and $L_2 = \mathcal{O}_W(a_2C_0 + b_2f)$, it follows that

$$\begin{aligned} a_1 + a_2 &= 3 \\ b_1 + b_2 &= m + e + 2 . \end{aligned}$$

We show now that $a_i \geq 1$ for $i = 1, 2$. Since φ is induced by the complete linear series of $\varphi^*\mathcal{O}_W(C_0 + mf)$, then $(1 - a_i)C_0 + (m - b_i)f$ is non effective. Then, if $a_i \leq 0$, $b_i \geq m + 1$. On the other hand, since X is regular and $H^1(L_i^*) \subset H^1(\varphi_*\mathcal{O}_X) = 0$, it follows that $H^1(\mathcal{O}_W(-a_iC_0 - b_if)) = 0$. Then, if $a_i \leq 0$, $b_i \leq 1$, hence $m \leq 0$, which is impossible, because $m \geq e + 1 \geq 1$. Then, since $a_1 + a_2 = 3$, a_i is either 1 or 2. Let us set $a_1 = 1$. Then, since $(1 - a_1)C_0 + (m - b_1)f$ cannot be effective, $b_1 \geq m + 1$. On the other hand, $a_2 = 2$, and

since $0 = h^1(L_2^*) = h^1(\mathcal{O}_W(-2C_0 - b_2f)) = h^1(\mathcal{O}_W((b_2 - e - 2)f)) = h^1(\mathcal{O}_{\mathbf{P}^1}(b_2 - e - 2))$, then $b_2 \geq e + 1$. Since $b_1 + b_2 = m + e + 2$, then $b_1 = m + 1$ and $b_2 = e + 1$. $\square$

The purpose of this paper is to study canonical quadruple Galois covers, and we will focus on them in the next sections. Meanwhile, Proposition 1.6 already yields a fact regarding these covers which is worth remarking. Note that a canonical Galois cover of the Veronese surface is flat, because the Veronese surface is smooth. Then, being φ flat and Galois, $\varphi_*\mathcal{O}_X$ splits completely, so we have this

Corollary 1.7. *There are no quadruple Galois canonical covers of the Veronese surface.*

2. QUADRUPLE GALOIS COVERS

In this section we start the study of the case when φ is a Galois, quadruple cover. We begin by setting up our definitions and recalling, without a proof, some useful facts on Galois extensions.

Definition 2.1. *Let $\mathfrak{X} \xrightarrow{\Phi} \mathfrak{W}$ be a finite morphism between normal projective varieties $\mathfrak{X}$ and $\mathfrak{W}$ and let G be a finite group acting on $\mathfrak{X}$ so that $\mathfrak{X}/G = \mathfrak{W}$ and Φ is in fact the projection from $\mathfrak{X}$ to $\mathfrak{X}/G$. We call $\mathfrak{X} \xrightarrow{\Phi} \mathfrak{W}$ a Galois cover with group G .*

Remark 2.2. *In the previous definition of Galois cover we are not assuming Φ to be flat. However, a quadruple Galois canonical cover $X \xrightarrow{\varphi} W$ of a smooth surface of minimal degree is flat, because W is smooth, X is locally Cohen-Macaulay and φ is finite.*

Lemma 2.3. *Let $A \subset B$ be a ring extension of integrally closed domains, let K be the field of fractions of A and let L be the field of fractions of B . Let G be a finite group. Then following are equivalent:*

- 1) G acts on B and $A = B^G$.
- 2) $K \subset L$ is a normal extension with Galois group G and B is the integral closure of A in L .

Under the equivalent conditions 1) and 2), G is the group of A -automorphisms of B .

Now we focus on Galois covers of degree 4. We recall the algebra structure associated to such covers:

Proposition 2.4. *Let $\mathfrak{X}$ and $\mathfrak{Y}$ be two algebraic varieties and let $\mathfrak{X} \xrightarrow{p} \mathfrak{Y}$ be a flat, Galois cover of degree 4.*

- 1) *If $G = \mathbf{Z}_4$, then $p_*\mathcal{O}_{\mathfrak{X}}$ splits as*

$$p_*\mathcal{O}_{\mathfrak{X}} = \mathcal{O}_{\mathfrak{Y}} \oplus L_i^* \oplus L_{-1}^* \oplus L_{-i}^*$$

where the line bundles on $\mathfrak{Y}$, $\mathcal{O}_{\mathfrak{Y}}, L_i^, L_{-1}^*$ and L_{-i}^* are the eigenspaces of $1, i, -1$ and $-i$ respectively.*

There exist effective Cartier divisors D_{11}, D_{12}, D_{23} and D_{33} on $\mathfrak{Y}$ such that $D_{11} + D_{23} = D_{12} + D_{33}$ and the following

$$\begin{aligned}
L_i \otimes L_i &= L_{-1} \otimes \mathcal{O}_Y(D_{11}) \\
L_i \otimes L_{-1} &= L_{-i} \otimes \mathcal{O}_Y(D_{12}) \\
L_i \otimes L_{-i} &= \mathcal{O}_Y(D_{11} + D_{23}) \\
L_{-1} \otimes L_{-1} &= \mathcal{O}_Y(D_{12} + D_{23}) \\
L_{-1} \otimes L_{-i} &= L_i \otimes \mathcal{O}_Y(D_{23}) \\
L_{-i} \otimes L_{-i} &= L_{-1} \otimes \mathcal{O}_Y(D_{33})
\end{aligned} \tag{2.4.1}$$

and the multiplicative structure of $p_*\mathcal{O}_{\mathfrak{X}}$ is as follows:

$$\begin{aligned}
L_i^* \otimes L_i^* &\xrightarrow{\cdot D_{11}} L_{-1}^* \\
L_i^* \otimes L_{-1}^* &\xrightarrow{\cdot D_{12}} L_{-i}^* \\
L_i^* \otimes L_{-i}^* &\xrightarrow{\cdot D_{11} + D_{23}} \mathcal{O}_{\mathfrak{Y}} \\
L_{-1}^* \otimes L_{-1}^* &\xrightarrow{\cdot D_{12} + D_{23}} \mathcal{O}_{\mathfrak{Y}} \\
L_{-1}^* \otimes L_{-i}^* &\xrightarrow{\cdot D_{23}} L_i^* \\
L_{-i}^* \otimes L_{-i}^* &\xrightarrow{\cdot D_{33}} L_{-1}^* .
\end{aligned}$$

2) If $G = \mathbf{Z}_2 \times \mathbf{Z}_2$, then $p_*\mathcal{O}_{\mathfrak{X}}$ splits as

$$p_*\mathcal{O}_{\mathfrak{X}} = \mathcal{O}_{\mathfrak{Y}} \oplus L_1^* \oplus L_2^* \oplus L_3^* ,$$

where $\mathcal{O}_{\mathfrak{Y}}, L_1^*, L_2^*$ and L_3^* are eigenspaces and there exist effective Cartier divisors D_1, D_2 and D_3 such that $L_i^{\otimes 2} = \mathcal{O}_Y(D_j + D_k)$ and $L_j \otimes L_k = L_i \otimes \mathcal{O}_Y(D_i)$ with $i \neq j, j \neq k$ and $k \neq i$, and the multiplicative structure of $p_*\mathcal{O}_{\mathfrak{X}}$ is as follows:

$$\begin{aligned}
L_i^* \otimes L_i^* &\xrightarrow{\cdot D_j + D_k} \mathcal{O}_{\mathfrak{Y}} , \\
L_j^* \otimes L_k^* &\xrightarrow{\cdot D_i} L_i^* .
\end{aligned}$$

Sketch of proof. The action of G splits $p_*\mathcal{O}_{\mathfrak{X}}$ into line bundles. Precisely if G is $\mathbf{Z}_4$, they are the eigenspaces of the eigenvalues $1, i, -1$ and $-i$ of one generator of the group and if G is $\mathbf{Z}_2 \times \mathbf{Z}_2$ they are the intersection of the eigenspaces associated to one of the generators of G with the eigenspaces of other generator of G . Then it is easy to trace to what eigenspace the product of elements of two eigenspaces goes. For instance, if G is $\mathbf{Z}_4$ it is clear that the product of two elements of L_i is an element of L_{-1} , the product of an element of L_i and an element of L_{-1} is an element of L_{-i} , etc. This yields the existence of the effective divisors mentioned in the statement and the description of the multiplicative structure, the relations among the divisors arising from the commutativity and associativity of the product. We refer to [Pa] and, for the bi-double case, to [C]; see also [HM]. $\square$

We recall also the definition of simple cyclic cover:

Definition-Remark 2.5. Let D be an effective Cartier divisor on a variety $\mathfrak{Y}$ such that there exists a line bundle L with $L^{\otimes n} = \mathcal{O}_{\mathfrak{Y}}(D)$. Consider the $\mathcal{O}_{\mathfrak{Y}}$ -algebra $\mathcal{O}_{\mathfrak{Y}} \oplus L^{-1} \oplus L^{-2} \oplus \dots \oplus L^{-n+1}$ with multiplication

$$\begin{aligned} L^{-m_1} \otimes L^{-m_2} &\longrightarrow L^{-(m_1+m_2)} \\ \text{if } 0 \leq m_1 + m_2 \leq n-1, &\quad \text{and} \\ L^{-m_1} \otimes L^{-m_2} &\xrightarrow{\cdot D} L^{-(m_1+m_2)+n} \\ &\text{otherwise.} \end{aligned}$$

Then we say that $\mathfrak{X} = \text{Spec}(\mathcal{O}_{\mathfrak{Y}} \oplus L^{-1} \oplus \dots \oplus L^{-n+1})$ is a simple cyclic cover of degree n branched along D .

A simple cyclic cover of degree n is Galois and its Galois group is $\mathbf{Z}_n$. In particular, with the notation of Proposition 2.4, a Galois cover $\mathfrak{X} \xrightarrow{p} \mathfrak{Y}$ with group $\mathbf{Z}_4$ is simple cyclic branched along D if and only if $D_{11} = D_{12} = 0$ and $D_{23} = D_{33} = D$ or $D_{11} = D_{12} = D$ and $D_{23} = D_{33} = 0$.

A quadruple Galois canonical cover over a smooth surface of minimal degree has the property that two of the eigenspaces (different from $\mathcal{O}_{\mathfrak{Y}}$) described in Proposition 2.4 have as tensor product the third one (see Proposition 1.3.) This is a strong condition that simplifies the algebra structures of the covers. We see what this exactly means in the next two propositions. We start with the case when the Galois group is $\mathbf{Z}_2 \times \mathbf{Z}_2$.

Proposition 2.6. Let $\mathfrak{X}$ and $\mathfrak{Y}$ be algebraic varieties and let $\mathfrak{X} \xrightarrow{p} \mathfrak{Y}$ be a flat Galois cover with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$. Let L_1, L_2, L_3, D_1, D_2 and D_3 be as in Proposition 2.4, 2). If $L_1 \otimes L_2 = L_3$, then

- 1) $\mathfrak{X}$ is the fiber product over $\mathfrak{Y}$ of the flat double covers $\mathfrak{X}_1 \xrightarrow{p_1} \mathfrak{Y}$ and $\mathfrak{X}_2 \xrightarrow{p_2} \mathfrak{Y}$, which are branched respectively along D_2 and D_1 and have trace zero module L_1^* and L_2^* respectively, and p is the natural map from the fiber product onto $\mathfrak{Y}$.
- 2) In particular, if $X \xrightarrow{\varphi} W$ is a canonical cover of a smooth surface of minimal degree and φ is Galois with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$, then X is the fiber product over W of two double covers $X_1 \xrightarrow{p_1} W$ and $X_2 \xrightarrow{p_2} W$ and φ is the natural map from the fiber product to W .

Proof. According to Proposition 2.4

$$p_*\mathcal{O}_{\mathfrak{X}} = \mathcal{O}_{\mathfrak{Y}} \oplus L_1^* \oplus L_2^* \oplus L_3^*$$

and the $\mathcal{O}_{\mathfrak{Y}}$ -algebra $p_*\mathcal{O}_{\mathfrak{X}}$ has three subalgebras $\mathcal{O}_{\mathfrak{Y}} \oplus L_1^*$, $\mathcal{O}_{\mathfrak{Y}} \oplus L_2^*$ and $\mathcal{O}_{\mathfrak{Y}} \oplus L_3^*$, corresponding to three double covers $\mathfrak{X}_i \xrightarrow{p_i} \mathfrak{Y}$ for $i = 1, 2, 3$ which are branched along $D_j + D_k$ where $i \neq j, j \neq k, k \neq i$. Since $L_1 \otimes L_2 = L_3$, then $D_3 = 0$, hence $\mathfrak{X}_1 \xrightarrow{p_1} \mathfrak{Y}$ is branched along D_2 and $\mathfrak{X}_2 \xrightarrow{p_2} \mathfrak{Y}$ is branched along D_1 . Then the algebra structure of $p_*\mathcal{O}_{\mathfrak{X}}$ described in Proposition 2.4, 2) becomes the tensor product over $\mathcal{O}_{\mathfrak{Y}}$ of the algebras $\mathcal{O}_{\mathfrak{Y}} \oplus L_1^*$ and $\mathcal{O}_{\mathfrak{Y}} \oplus L_2^*$.

Now let $X \xrightarrow{\varphi} W$ be a canonical cover of degree 4. If φ is Galois, by Proposition 2.4, $\varphi_*\mathcal{O}_X$ splits as direct sum of line bundles. Then by Proposition 1.3 $L_1 \otimes L_2 = L_3$, so the statement follows from the first part of Proposition 2.6. $\square$

Proposition 2.7. *Let $\mathfrak{X}$ and $\mathfrak{Y}$ be algebraic varieties and let $\mathfrak{X} \xrightarrow{p} \mathfrak{Y}$ be a flat Galois cover with Galois group $\mathbf{Z}_4$ and let $L_i, L_{-1}, L_{-i}, D_{11}, D_{12}, D_{23}$ and D_{33} be as in Proposition 2.4. If $L_{\tau(i)} \otimes L_{\tau(-1)} = L_{\tau(-i)}$ for some permutation τ of $\{i, -1, -i\}$, then*

- 1) *either $L_i \otimes L_{-1} = L_{-i}$ (i.e., $D_{12} = 0$) or $L_{-1} \otimes L_{-i} = L_i$ (i.e., $D_{23} = 0$).*
- 2) *In the case $L_i \otimes L_{-1} = L_{-i}$, the cover $\mathfrak{X} \xrightarrow{p} \mathfrak{Y}$ is the composition of a flat double cover $\mathfrak{X}' \xrightarrow{p_1} \mathfrak{Y}$ branched along D_{23} followed by a flat double cover $\mathfrak{X} \xrightarrow{p_2} \mathfrak{X}'$, branched along $p_1^* D_{11}$ and the ramification locus of p_1 . Moreover, the trace zero module of p_2 is $p_1^* L_i^*$ and the trace zero module of p_1 is L_{-1}^* . The surface $\mathfrak{X}'$ is the quotient of $\mathfrak{X}$ by the unique subgroup of index 2 of $\mathbf{Z}_4$. If $L_{-1} \otimes L_{-i} = L_i$ there exists an analogous decomposition of p .*
- 3) *Assume that $\mathfrak{Y}$ is locally Gorenstein. If $L_i \otimes L_{-1} = L_{-i}$, then $\mathfrak{X}$ is locally Gorenstein and $\omega_{\mathfrak{X}} = p^*(\omega_{\mathfrak{Y}} \otimes L_{-i})$. If $L_{-1} \otimes L_{-i} = L_i$, then $\mathfrak{X}$ is locally Gorenstein and $\omega_{\mathfrak{X}} = p^*(\omega_{\mathfrak{Y}} \otimes L_i)$.*
- 4) *In particular, if $X \xrightarrow{\varphi} W$ is a canonical cover of degree 4 of a smooth surface of minimal degree and φ is Galois with Galois group $\mathbf{Z}_4$, then $X \xrightarrow{\varphi} W$ is a cover satisfying 1), 2) and 3) above and either L_i or L_{-i} is isomorphic to $\omega_W^*(1)$.*

Proof. By assumption either

$$\begin{aligned} L_i \otimes L_{-1} &= L_{-i} \text{ or} \\ L_i \otimes L_{-i} &= L_{-1} \text{ or} \\ L_{-1} \otimes L_{-i} &= L_i. \end{aligned}$$

Since $\mathfrak{X}$ is normal, $L_i \otimes L_{-i} = L_{-1}$ is not possible. Then either $L_i \otimes L_{-1} = L_{-i}$ or $L_{-1} \otimes L_{-i} = L_i$ so in Proposition 2.4 either $D_{12} = 0$ or $D_{23} = 0$. If, for example, $D_{12} = 0$, then looking locally at the multiplicative structure of $p_* \mathcal{O}_X$ yields the description of the trace zero module and branch loci given in 2). Now, L_{-1}^* is the trace zero module of p_1 and $p_1^* L_i^*$ is the trace zero module of p_2 , so $\omega_{\mathfrak{X}} = p^*(\omega_{\mathfrak{Y}} \otimes L_i \otimes L_{-1}) = p^*(\omega_{\mathfrak{Y}} \otimes L_{-i})$. This proves 3). Finally, from Proposition 1.3 we know also that, if $X \xrightarrow{\varphi} W$ is a canonical cover of degree 4, then $L_{\tau(i)} \otimes L_{\tau(-1)} = L_{\tau(-i)}$ for some permutation τ of $\{i, -1, -i\}$, so $X \xrightarrow{\varphi} W$ satisfies the hypothesis of the statement. Then by 1) either $L_i \otimes L_{-1} = L_{-i}$ or $L_{-1} \otimes L_{-i} = L_i$. We argue for instance in case $L_i \otimes L_{-1} = L_{-i}$. Since $\omega_X = p^* \omega_W(1)$, then 3) implies that L_{-i} is numerically equivalent to $\omega_W^*(1)$, but since W is either $\mathbf{P}^2$ or a Hirzebruch surface we have $L_{-i} = \omega_W^*(1)$. If $L_{-1} \otimes L_{-i} = L_i$ arguing similarly we obtain $L_i = \omega_W^*(1)$. $\square$

Notation-Remark 2.8. *If $X \xrightarrow{\varphi} W$ is a canonical cover of degree 4 of a smooth surface of minimal degree and φ is Galois with group $\mathbf{Z}_4$, then we rename $\{L_i, L_{-1}, L_{-i}\}$ as $\{L_1, L_2, L_3\}$ so that $L_1 \otimes L_2 = L_3$, $L_2 = L_{-1}$ and $L_3 = \omega_W^*(1)$. Then there exist effective*

Cartier divisors D_1 and D_2 so that

$$\begin{aligned} L_1 \otimes L_1 &= L_2 \otimes \mathcal{O}_W(D_1) \\ L_1 \otimes L_2 &= L_3 \\ L_1 \otimes L_3 &= \mathcal{O}_W(D_1 + D_2) \\ L_2 \otimes L_2 &= \mathcal{O}_W(D_2) \\ L_2 \otimes L_3 &= L_1 \otimes \mathcal{O}_W(D_2) \\ L_3 \otimes L_3 &= L_2 \otimes \mathcal{O}_W(D_1 + D_2) . \end{aligned}$$

This is achieved if we set $D_1 = D_{11}$ and $D_2 = D_{23}$ in (2.4.1) in case $L_1 = L_i$ and one can argue analogously if $L_1 = L_{-i}$. In particular φ is simple cyclic if and only if $D_1 = 0$.

Now we make two useful observations which are a sort of converses to Proposition 2.6 and Proposition 2.7:

Proposition 2.9. *Let $\mathfrak{X}$ and $\mathfrak{Y}$ be normal algebraic varieties.*

- 1) *If $\mathfrak{X} \xrightarrow{p} \mathfrak{Y}$ is a morphism such that p is the natural map onto $\mathfrak{Y}$ from the fiber product over $\mathfrak{Y}$ of two flat double covers $\mathfrak{X}_1 \xrightarrow{p_1} \mathfrak{Y}$ and $\mathfrak{X}_2 \xrightarrow{p_2} \mathfrak{Y}$, then p is a Galois cover with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$.*
- 2) *If in addition L_2^* and L_1^* are the trace zero modules of p_2 and p_1 respectively, then*

$$p_*\mathcal{O}_{\mathfrak{X}} = \mathcal{O}_{\mathfrak{Y}} \oplus L_1^* \oplus L_2^* \oplus (L_1^* \otimes L_2^*)$$

and, if $\mathfrak{Y}$ is locally Gorenstein, then $\mathfrak{X}$ is locally Gorenstein and $\omega_{\mathfrak{X}} = p^(\omega_{\mathfrak{Y}} \otimes L_1 \otimes L_2)$.*

Proof. Choosing suitable open sets U and V of $\mathfrak{X}$ and $\mathfrak{Y}$ we can describe the structure of $\mathcal{O}_{\mathfrak{Y}}$ -algebra of $\mathcal{O}_{\mathfrak{X}}$ in local coordinates and see that $\mathbf{Z}_2 \oplus \mathbf{Z}_2$ acts on $\mathcal{O}_U$ and its ring of invariants is $\mathcal{O}_V$. Then, since $\mathfrak{X}$ and $\mathfrak{Y}$ are normal, by Lemma 2.3 the extension $\mathcal{K}(X)/\mathcal{K}(Y)$ is Galois with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$, and since $\mathfrak{X}$ and $\mathfrak{Y}$ are normal and p is finite, $\mathcal{O}_{\mathfrak{X}}$ is the integral closure of $\mathcal{O}_{\mathfrak{Y}}$ in $\mathcal{K}(X)$, so again by Lemma 2.3 the morphism p is a Galois cover with group $\mathbf{Z}_2 \times \mathbf{Z}_2$. The computation of the push-forward of $\mathcal{O}_{\mathfrak{X}}$ and the computation of the canonical of $\mathfrak{X}$ is straight-forward once we recall that the trace zero module of the double cover p_1 is L_1^* and using the fact that $p = p_1 \circ p'_2$, where p'_2 is another double cover with trace zero module $p_1 L_2^*$. $\square$

Proposition 2.10. *Let $\mathfrak{X}$ and $\mathfrak{Y}$ be normal algebraic varieties.*

- 1) *If $\mathfrak{X} \xrightarrow{p} \mathfrak{Y}$ is the composition of a flat double cover $\mathfrak{X}' \xrightarrow{p_1} \mathfrak{Y}$ branched along a divisor D_2 , followed by a flat double cover $\mathfrak{X} \xrightarrow{p_2} \mathfrak{X}'$, branched along the ramification locus of p_1 and $p_1^* D_1$, where D_1 is a divisor on $\mathfrak{Y}$, then p is a Galois cover with Galois group $\mathbf{Z}_4$.*
- 2) *If in addition L_2^* is the trace zero module of p_1 and $p_1^* L_1^*$ is the trace zero module of p_2 , then*

$$p_*\mathcal{O}_{\mathfrak{X}} = \mathcal{O}_{\mathfrak{Y}} \oplus L_1^* \oplus L_2^* \oplus (L_1^* \otimes L_2^*)$$

and, if $\mathfrak{Y}$ is locally Gorenstein, then $\mathfrak{X}$ is locally Gorenstein and $\omega_{\mathfrak{X}} = p^(\omega_{\mathfrak{Y}} \otimes L_1 \otimes L_2)$.*

Proof. Let V a smooth open set of $\mathfrak{Y}$ contained in $\mathfrak{Y} - D_1$. Over V the cover p is a simple cyclic cover, and in particular, a Galois cover with Galois group $\mathbf{Z}_4$. Arguing as in Proposition 2.9 we conclude that $\mathfrak{X} \xrightarrow{p} \mathfrak{Y}$ is Galois with Galois group $\mathbf{Z}_4$. The computation of the push-forward of $\mathcal{O}_{\mathfrak{X}}$ and of the canonical bundle of $\mathfrak{X}$ is straight-forward from the fact that $p = p_1 \circ p_2$ and the knowledge of the trace zero modules of p_1 and p_2 . $\square$

In the remaining of the paper we will classify quadruple Galois canonical covers of smooth surfaces W of minimal degree. A priori one could distinguish three cases: W is linear $\mathbf{P}^2$, W is the Veronese surface and W is a smooth rational normal scroll. However, as pointed out in Corollary 1.7, there are no canonical quadruple Galois covers of the Veronese surface, so we will have only to study the cases of W being linear $\mathbf{P}^2$, dealt with in the next theorem, and of W being a smooth rational normal scroll, dealt with in Section 3.

Theorem 2.11. *Let W be linear $\mathbf{P}^2$ and let $X \xrightarrow{\varphi} W$ be a canonical Galois cover of degree 4.*

- 1) *If the Galois group of φ is $\mathbf{Z}_4$, then φ is the composition of two flat double covers $X_1 \xrightarrow{p_1} W$ and $X \xrightarrow{p_2} X_1$; the cover p_1 is branched along a quartic and the cover p_2 is branched along the ramification of p_1 and the pullback by p_1 of a conic and its trace zero module is $p_1^* \mathcal{O}_{\mathbf{P}^2}(2)$.*
- 2) *If the Galois group of φ is $\mathbf{Z}_2 \times \mathbf{Z}_2$ then X is the fiber product over W of two double covers of linear $\mathbf{P}^2$, each of them branched along a quartic, and φ is the natural map from the fiber product to W .*

Conversely, let $X \xrightarrow{\varphi} W$ be a cover of linear $\mathbf{P}^2$.

- 1') *If φ is the composition of two flat double covers $X_1 \xrightarrow{p_1} W$ and $X \xrightarrow{p_2} X_1$ as described in 1) above, then φ is a Galois canonical cover with group $\mathbf{Z}_4$.*
- 2') *If φ is the natural map to W from the fiber product over W of two double covers as described in 2) above, then φ is a Galois canonical cover with group $\mathbf{Z}_2 \times \mathbf{Z}_2$.*

Proof. By Proposition 2.4 the bundle $\varphi_* \mathcal{O}_X$ splits as a sum of line bundles. Now 1) and 2) follow from Proposition 1.6, 1), Proposition 2.6 and Proposition 2.7. Now we prove the converse. If φ is a cover as in 1'), then by Proposition 2.10 it is Galois with Galois group $\mathbf{Z}_4$. Likewise if φ is a cover as in 2'), then by Proposition 2.9 it is Galois with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$. On the other hand, by Proposition 2.9 and Proposition 2.10 the canonical of X is $\varphi^* \mathcal{O}_{\mathbf{P}^2}(1)$, so X is a surface of general type and ω_X is base-point-free. Finally to prove that φ is indeed the morphism induced by $H^0(\omega_X)$ we compare $H^0(\varphi^* \mathcal{O}_{\mathbf{P}^2}(1))$ and $H^0(\mathcal{O}_{\mathbf{P}^2}(1))$ and see that they are equal. Indeed, to compute $H^0(\varphi^* \mathcal{O}_{\mathbf{P}^2}(1))$ we push down $\varphi^* \mathcal{O}_{\mathbf{P}^2}(1)$ to W , and since

$$\varphi_* \mathcal{O}_X = \mathcal{O}_{\mathbf{P}^2} \oplus \mathcal{O}_{\mathbf{P}^2}(-2) \oplus \mathcal{O}_{\mathbf{P}^2}(-2) \oplus \mathcal{O}_{\mathbf{P}^2}(-4)$$

by Proposition 2.9 and Proposition 2.10, we obtain $H^0(\varphi^* \mathcal{O}_{\mathbf{P}^2}(1)) = H^0(\mathcal{O}_{\mathbf{P}^2}(1))$ as wished. $\square$

We describe further the Galois covers appearing in Theorem 2.11:

Corollary 2.12. *Let W be linear $\mathbf{P}^2$ and let $X \xrightarrow{\varphi} W$ be a Galois canonical cover of degree 4. Then,*

1) *the push-forward of $\mathcal{O}_X$ is as follows:*

$$\varphi_*\mathcal{O}_X = \mathcal{O}_{\mathbf{P}^2} \oplus \mathcal{O}_{\mathbf{P}^2}(-2) \oplus \mathcal{O}_{\mathbf{P}^2}(-2) \oplus \mathcal{O}_{\mathbf{P}^2}(-4) ;$$

2) *the surface X is regular; and*

3) *if the Galois group of φ is $\mathbf{Z}_4$, then X is singular and the mildest possible set of singularities on X consists of 8 points of type A_1 .*

Proof. Claim 1) of the corollary has already been shown in the proof of Theorem 2.11. Claim 2) is straight-forward from 1), since $H^1(\mathcal{O}_X) = H^1(\varphi^*\mathcal{O}_X)$. To prove 3) just observe that $X_1 \xrightarrow{p_1} W$ is branched along a quartic D_2 of $\mathbf{P}^2$ and p_2 along the ramification of p_1 and $p_1^*D_1$, where D_1 is a conic of $\mathbf{P}^2$. If D_1 and D_2 are both smooth and meet transversally then X_1 is smooth and the branch locus of p_2 has 8 singular points of type A_1 so X is smooth except at 8 points, which are singularities of type A_1 . $\square$

We end the section by remarking the existence of examples of covers like those appearing in Theorem 2.11:

Proposition 2.13. *Let W be linear $\mathbf{P}^2$.*

- 1) *There exist canonical covers $X \xrightarrow{\varphi} W$ with Galois group $\mathbf{Z}_4$ (that is, covers as in Theorem 2.11, 1)) with 8 singularities of type A_1 as only singularities.*
- 2) *There exist canonical covers $X \xrightarrow{\varphi} W$ with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$ (that is, covers as in Theorem 2.11, 2)) with X smooth.*

Proof. We first deal with 1). By the converse part in Theorem 2.11 we just have to construct a composition of double covers $X \xrightarrow{p_2} X_1$ and $X_1 \xrightarrow{p_1} W$ branched along suitable divisors. Then, using the same notation of the proof of Corollary 2.12 and by the argument there, it suffices to choose D_1 a smooth quartic and D_2 a smooth conic of $\mathbf{P}^2$ meeting transversally. This is possible by Bertini. Now for 2) again by the converse part in Theorem 2.11 we just have to construct the fiber product φ of two double covers $X_1 \xrightarrow{p_1} \mathbf{P}^2$ and $X_2 \xrightarrow{p_2} \mathbf{P}^2$ branched along suitable divisors D_2 and D_1 , which are two quartics in this case. Then $\varphi = p_1 \circ p_2'$ where $X \xrightarrow{p_2'} X_1$ is a double cover branched along $p_1^*D_1$. If we choose D_1 and D_2 smooth and meeting transversally (which again is possible by Bertini), then X_1 is smooth and so is the branch locus of p_2' , hence X is also smooth. Note that one can construct examples of X with worse singularities by allowing $D_1 + D_2$ to have worse singularities. $\square$

3. GALOIS COVERS OF SMOOTH RATIONAL NORMAL SCROLLS

At the end of Section 2 we saw that quadruple Galois canonical covers of $\mathbf{P}^2$ can have Galois group $\mathbf{Z}_4$ or $\mathbf{Z}_2 \times \mathbf{Z}_2$, but, in the first case, they could not be simple cyclic. The same thing happens for the covers we study in this section and we start by showing that there do not exist simple cyclic quadruple canonical covers X of smooth rational normal

scrolls. This was known if X is regular and W is a surface of minimal degree, whether smooth or singular, as consequence of the non-existence of canonical simple cyclic covers of degree bigger than 3 (see [GP2]). In the next theorem we prove the non existence of simple cyclic covers of degree bigger than 3 when X is an arbitrary surface of general type and W is a smooth rational scroll.

Theorem 3.1. *Let W be a smooth rational normal scroll, let $X \xrightarrow{\varphi} W$ be a canonical cover of degree n . If $X \xrightarrow{\varphi} W$ is a Galois cover and $n \geq 4$, then φ is not simple cyclic.*

Proof. Let us assume φ is simple cyclic. Then

$$\varphi_* \mathcal{O}_X = \mathcal{O}_W \oplus L^{-1} \oplus \dots \oplus L^{-n+1}.$$

Recall that W is isomorphic to $\mathbf{F}_e$. On the one hand $\omega_X = \varphi^* \mathcal{O}_W(1) = \varphi^* \mathcal{O}_W(C_0 + mf)$, with $m \geq e + 1$. On the other hand, $\omega_X = \varphi^*(\omega_W \otimes L^{-n+1}) = \varphi^*(\mathcal{O}_W(-2C_0 - (e+2)f) \otimes L^{-n+1})$. Thus $\varphi^* L^{-n+1} = \varphi^* \mathcal{O}_W(3C_0 + (m+e+2)f)$, so L^{-n+1} and $\mathcal{O}_W(3C_0 + (m+e+2)f)$ are numerically equivalent in W . Since W is a Hirzebruch surface, $L^{-n+1} = \mathcal{O}_W(3C_0 + (m+e+2)f)$ and 3 and $m+e+2$ are both multiple of $n-1$. Since $n \geq 4$ by assumption, this makes $n = 4$. In that case, since φ is induced by the complete series of $\mathcal{O}_W(C_0 + mf)$, then $\mathcal{O}_W((m - \frac{1}{3}(m+e+2))f)$ should be non effective, i.e., $m - \frac{1}{3}(m+e+2) < 0$, which is the same as $2m - e - 2 < 0$. Then, since $m \geq e + 1$, we get the contradiction $e < 0$ and φ cannot be simple cyclic. $\square$

After Theorem 3.1 we summarize now the status of the existence of simple cyclic canonical covers in the following theorem. To see the scope of the result, we remark that Theorem 3.2 implies the non-existence of Galois canonical covers of prime degree p of smooth scrolls, $\mathbf{P}^2$ or the Veronese surface, if $p \geq 5$. If in addition X is regular, next theorem assures that, if $p \geq 5$, then there are no Galois canonical covers of prime degree p of any surface of minimal degree.

Theorem 3.2. *Let W be a surface of minimal degree, not necessarily smooth, and let $X \xrightarrow{\varphi} W$ be a Galois canonical cover. If X is regular or W is smooth, and if φ is simple cyclic, then $\deg \varphi \leq 3$.*

Proof. If X is regular, the result follows from [GP2], Corollary 3.2. So we will assume that X is irregular and W smooth. The surface W cannot be isomorphic to $\mathbf{P}^2$, for if it were, since φ is simple cyclic, $\varphi_* \mathcal{O}_X$ would split completely, and so X would be regular. Thus W is a smooth rational normal scroll. Then we conclude that $\deg \varphi \leq 3$ by applying Theorem 3.1. $\square$

Corollary 3.3. *Let W be a surface of minimal degree, not necessarily smooth, and let $X \xrightarrow{\varphi} W$ be a Galois canonical cover. If X is regular or W is smooth and φ is a Galois canonical cover of prime degree, then $\deg \varphi \leq 3$.*

Proof. If $\deg \varphi = p$ is prime, then G is cyclic of order p , so the stabilizer of any $x \in X$ is either $\{id\}$ or G , so φ is simple cyclic. $\square$

These results hint towards a positive solution to the following very interesting question regarding Galois canonical covers of prime degree bigger than 3.

Question 3.4. *If $X \xrightarrow{\varphi} W$ is a Galois canonical cover of prime degree, is then $\deg \varphi \leq 3$?*

Theorem 3.1 showed that there does not exist simple cyclic canonical covers of degree bigger than 3 of smooth rational normal scrolls. The construction of non-simple cyclic canonical covers of surfaces is not easy and there are not many examples, to the best of the authors knowledge. We now prove a classification theorem for non-simple cyclic canonical covers $X \xrightarrow{\varphi} W$ of degree 4, splitting the cases X regular and X irregular. Theorem 3.5 and Theorem 3.6 below and Proposition 3.11 show that they do exist, but, as Corollary 3.7 tells, they are always singular having at best singularities of type A_1 .

Theorem 3.5. *Let $X \xrightarrow{\varphi} W$ be a Galois canonical cover of degree 4 and Galois group $\mathbf{Z}_4$ with X regular and let $W = S(m - e, m)$ be a smooth rational normal scroll. Then φ is the composition of two flat double covers $X_1 \xrightarrow{p_1} W$ and $X \xrightarrow{p_2} X_1$ which are as follows:*

The cover p_1 is branched along a divisor D_2 on W .

*The cover p_2 is branched along the ramification of p_1 and $p_1^*D_1$ and has trace zero module $p_1^*\mathcal{O}_Y(-\frac{1}{2}D_1 - \frac{1}{4}D_2)$, where D_1 is a divisor on W .*

The scroll W and the divisors D_1 and D_2 satisfy:

- 1) $W = S(m - e, m)$, $0 \leq e \leq 2$, $m \geq e + 1$, $D_1 \sim (2m - e + 1)f$, $D_2 \sim 4C_0 + (2e + 2)f$
- 2) $W = S(1, 1)$ (i.e., W is a smooth quadric hypersurface in $\mathbf{P}^3$), $D_1 \sim 3C_0$ and $D_2 \sim 2C_0 + 4f$

Conversely, let $X \xrightarrow{\varphi} W$ be the composition of two flat double covers $X_1 \xrightarrow{p_1} W$ and $X \xrightarrow{p_2} X_1$ as described above, then φ is a Galois canonical cover of W with Galois group $\mathbf{Z}_4$ and X is regular.

Proof. Proposition 2.7, 4) says that φ is the composition of two double covers $X_1 \xrightarrow{p_1} W$ branched along a divisor D_2 and $X \xrightarrow{p_2} X_1$, branched along the ramification of p_1 and $p_1^*D_1$ and, according to Notation-Remark 2.8,

$$\varphi_*\mathcal{O}_X = \mathcal{O}_W \oplus L_1^* \oplus L_2^* \oplus L_3^*,$$

where L_2^* is the eigenspace for -1 , L_1^* is the eigenspace for either i or $-i$, L_3^* the eigenspace for $-i$ or i and $L_1 \otimes L_2 = L_3$. Moreover, $L_3 = \omega_W^*(1)$, $L_1^{\otimes 2} = L_2 \otimes \mathcal{O}_W(D_1)$, $L_2^{\otimes 2} = \mathcal{O}_W(D_2)$, $p_1^*L_1^*$ is the trace zero module of p_2 and L_2^* is the trace zero module of p_1 . From this we obtain that the trace zero module of p_2 is $p_1^*L_1^* = p_1^*\mathcal{O}_Y(-\frac{1}{2}D_1 - \frac{1}{4}D_2)$.

Now we show that W, D_1 and D_2 satisfy 1) or 2). Recall that W is isomorphic to $\mathbf{F}_e$. Since X is regular we can apply Proposition 1.6, 3). Then, since $L_3 = \omega_W^*(1)$ we have either $L_1^* = \mathcal{O}_W(-C_0 - (m+1)f)$, $L_2^* = \mathcal{O}_W(-2C_0 - (e+1)f)$ and $L_3^* = \mathcal{O}_W(-3C_0 - (m+e+2)f)$ or $L_2^* = \mathcal{O}_W(-C_0 - (m+1)f)$, $L_1^* = \mathcal{O}_W(-2C_0 - (e+1)f)$ and $L_3^* = \mathcal{O}_W(-3C_0 - (m+e+2)f)$

Case 1: $L_1^* = \mathcal{O}_W(-C_0 - (m+1)f)$, $L_2^* = \mathcal{O}_W(-2C_0 - (e+1)f)$. From the previous description, φ is the composition of $X_1 \xrightarrow{p_1} W$, where p_1 is a double cover branched along a divisor D_2 linearly equivalent to $4C_0 + (2e+2)f$ and $X \xrightarrow{p_2} X_1$, where p_2 is a double cover branched along the ramification of p_1 and $p_1^*D_1$, where D_1 is linearly equivalent to

$(2m - e + 1)f$. Recall that X is normal, hence the components of $D_1 + D_2$ have multiplicity 1 and in particular, the fixed part of $|4C_0 + (2e + 2)f|$ contains C_0 with multiplicity at most 1. Thus $e \leq 2$.

Case 2: $L_2^* = \mathcal{O}_W(-C_0 - (m + 1)f)$, $L_1^* = \mathcal{O}_W(-2C_0 - (e + 1)f)$. Again from the description above, φ is the composition of $X_1 \xrightarrow{p_1} W$, where p_1 is a double cover branched along a divisor D_2 linearly equivalent to $2C_0 + (2m + 2)f$ and $X \xrightarrow{p_2} X_1$, where p_2 is a double cover branched along the ramification of p_1 and $p_1^*D_1$, where D_1 is linearly equivalent to $3C_0 + (2e - m + 1)f$. Recall that X is normal, hence the components of $D_1 + D_2$ have multiplicity 1 and, in particular, the fixed part of $|3C_0 + (2e - m + 1)f|$ contains C_0 with multiplicity at most 1. Thus $m \leq 1$, and in fact $m = 1$ and $e = 0$. So finally, $W = \mathbf{F}_0$, D_1 is linearly equivalent to $3C_0$ and D_2 is linearly equivalent to $2C_0 + 4f$.

Now we prove the converse. By Proposition 2.10 the morphism φ is a Galois cover with group $\mathbf{Z}_4$. Proposition 2.10 also tells us that $\omega_X = \varphi^*(\omega_W \otimes L_1 \otimes L_2)$. Since $L_1 \otimes L_2 = \mathcal{O}_W(3C_0 + (m + e + 2)f) = \omega_W^*(1)$ in either 1) or 2), we have $\omega_X = \varphi^*\mathcal{O}_W(1)$, so X is a surface of general type with base-point-free canonical bundle. Now, again by Proposition 2.10, in both 1) and 2) we have

$$\begin{aligned} \varphi_*\mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-C_0 - (m + 1)f) \oplus \\ &\mathcal{O}_W(-2C_0 - (e + 1)f) \oplus \mathcal{O}_W(-3C_0 - (m + e + 2)f). \end{aligned} \quad (3.5.1)$$

Then to see that φ is the canonical morphism of X we compare $H^0(\omega_X) = H^0(\varphi^*\mathcal{O}_W(1))$ and $H^0(\mathcal{O}_W(1))$. The group $H^0(\varphi^*\mathcal{O}_W(1))$ can be computed pushing $\varphi^*\mathcal{O}_W(1)$ down to W and using (3.5.1), and one sees at once that $H^0(\omega_X) = H^0(\mathcal{O}_W(1))$. Finally we see that $H^1(\mathcal{O}_X) = 0$ also by pushing down to W and using (3.5.1). $\square$

Theorem 3.6. *Let $X \xrightarrow{\varphi} W$ be a canonical Galois cover of degree 4 and Galois group $\mathbf{Z}_4$ with X irregular and let $W = S(m - e, m)$ be a smooth rational normal scroll. Then the irregularity of X is $q(X) = 1$ and W is isomorphic to $\mathbf{F}_0$. Moreover, φ is the composition of two flat double covers $X_1 \xrightarrow{p_1} W$ and $X \xrightarrow{p_2} X_1$ which are as follows:*

The cover p_1 is branched along a divisor D_2 on W .

*The cover p_2 is branched along the ramification of p_1 and $p_1^*D_1$ and has trace zero module $p_1^*\mathcal{O}_W(-\frac{1}{2}D_1 - \frac{1}{4}D_2)$, where D_1 is a divisor on W .*

The scroll W and the divisors D_1 and D_2 satisfy:

- 1) $W = S(m, m)$, $m \geq 1$, $D_1 \sim (2m + 4)f$ and $D_2 \sim 4C_0$.
- 2) $W = S(1, 1)$, $D_1 \sim 6C_0$ and $D_2 \sim 4f$.

Conversely, if $X \xrightarrow{\varphi} W$ is the composition of two double covers $X_1 \xrightarrow{p_1} W$ and $X \xrightarrow{p_2} X_1$ as described above, then φ is a canonical Galois cover of W with Galois group $\mathbf{Z}_4$ and X is irregular.

Proof. Using Proposition 2.7, 4) and Notation-Remark 2.8 as in the proof of Theorem 3.5 we conclude that φ is the composition of two double covers. The first one is $X_1 \xrightarrow{p_1} W$, is branched along a divisor D_2 and has trace zero module L_2^* . The second cover is $X \xrightarrow{p_2} X_1$,

is branched along the ramification of p_1 and $p_1^*D_1$ and has trace zero module $p_1^*L_1$. Then $L_1^{\otimes 2} = L_2 \otimes \mathcal{O}_W(D_1)$, $L_2^{\otimes 2} = \mathcal{O}_W(D_2)$ and moreover, $L_1 \otimes L_2 = L_3 = \omega_W^*(1)$. Recall that $W = \mathbf{F}_e$ and let $L_1 = \mathcal{O}_W(a_1C_0 + b_1f)$ and $L_2 = \mathcal{O}_W(a_2C_0 + b_2f)$. Then we have

$$\begin{aligned} a_1 + a_2 &= 3 \\ b_1 + b_2 &= m + e + 2 \end{aligned}$$

Since $L_1^{\otimes 2} \otimes L_2^* = \mathcal{O}_W(D_1)$ and $L_2^{\otimes 2} = \mathcal{O}_W(D_2)$ are effective then $a_2 \leq 2a_1$, $b_2 \leq 2b_1$ and $a_2, b_2 \geq 0$. Then $a_1, b_1 \geq 0$ also. Moreover, $a_1, b_1 \geq 1$, otherwise we will contradict $a_1 + a_2 = 3$ or $b_1 + b_2 = m + e + 2$. We see that a_1 cannot be 2. If $a_1 = 2$, then $D_1 \sim 3C_0 + (2b_1 - b_2)f$, $D_2 \sim 2C_0 + 2b_2f$, $L_1 = \mathcal{O}_W(2C_0 + b_1f)$ and $L_2 = \mathcal{O}_W(C_0 + b_2f)$. Since X is irregular, and $H^1(\mathcal{O}_W) = H^1(L_2^*) = H^1(L_3^*) = 0$, then $H^1(L_1^*) \neq 0$. This implies $b_1 \leq e$. Then $b_1 + b_2 = m + e + 2$ implies $b_2 \geq m + 2$. On the other hand, since X is normal, C_0 has at most multiplicity 1 in the fixed part of $|3C_0 + (2b_1 - b_2)f|$, and this implies $2b_1 - b_2 - 2e \geq 0$. Then we have $2e - (m + 2) - 2e \geq 0$, which is a contradiction. Then the only possibilities are $a_1 = 1$ or $a_1 = 3$.

Case 1: $a_1 = 1$. Then $a_2 = 2$ and X irregular implies $b_2 \leq e$, since L_2^* has to be special. The fact that X is normal implies $-3e + 2b_2 \geq 0$, since D_2 cannot have $2C_0$ as a fixed component. Then $b_2 = e = 0$, and, summarizing, $e = 0, a_1 = 1, a_2 = 2, b_1 = m + 2, b_2 = 0$. This implies $D_1 \sim (2m + 4)f$ and $D_2 \sim 4C_0$.

Case 2: $a_1 = 3$. Then $D_1 \sim 6C_0 + (2b_1 - b_2)f$, and since X is normal, C_0 has at most multiplicity 1 in the fixed part of $|D_1|$, hence $2b_1 - b_2 - 5e \geq 0$. Now since $H^0(\varphi^*\mathcal{O}_W(C_0 + mf)) = H^0(\mathcal{O}_W(C_0 + mf))$, we have that $b_2 > m$, hence $b_1 < e + 2$. Then we get $2e + 4 - m - 5e = -3e - m + 4 > 0$. But $m \geq e + 1$, so this gives $-4e + 3 > 0$, hence $e = 0$. In this case, $b_1 = 1$ and $m \geq 1$. Then $b_2 = m + 1 \geq 2$, and since D_1 is effective, $b_2 = 2$ and $m = 1$. Summarizing, $e = 0, m = 1, a_1 = 3, a_2 = 0, b_1 = 1$ and $b_2 = 2$. Then $D_1 \sim 6C_0$ and $D_2 \sim 4f$.

Finally the irregularity of X is $h^1(\mathcal{O}_X) = h^1(\varphi_*\mathcal{O}_X)$. We observe that the computation of a_1, a_2, b_1 and b_2 yields that $\varphi_*\mathcal{O}_X$ is

$$\begin{aligned} &\mathcal{O}_W \oplus \mathcal{O}_W(-C_0 - (m + 2)f) \oplus \mathcal{O}_W(-2C_0) \oplus \mathcal{O}_W(-3C_0 - (m + 2)f) \text{ or} \\ &\mathcal{O}_W \oplus \mathcal{O}_W(-3C_0 - f) \oplus \mathcal{O}_W(-2f) \oplus \mathcal{O}_W(-3C_0 - 3f) \quad (3.6.1) \end{aligned}$$

so $h^1(\mathcal{O}_X) = h^1(\mathcal{O}_W(-2C_0))$ or $h^1(\mathcal{O}_X) = h^1(\mathcal{O}_W(-2f))$, and in both cases, equal to 1.

We prove now the converse. Proposition 2.10 implies that φ is Galois with Galois group $\mathbf{Z}_4$. Now if L_2^* is the trace zero module of p_1 and $p_1^*L_1^*$ is the trace zero module of p_2 , then $L_1 \otimes L_2 = \mathcal{O}_W(\frac{1}{2}D_1 + \frac{3}{4}D_2)$. Then if D_1 and D_2 are as in 1) or 2), $L_1 \otimes L_2 = \omega_W^*(1)$. Then Proposition 2.10 implies also that $\omega_X = \varphi^*\mathcal{O}_W(1)$, therefore X is a surface of general type with base-point-free canonical bundle. Finally Proposition 2.10 tells us also that $\varphi_*\mathcal{O}_X$ is as in (3.6.1) so arguing like in the end of the proof of Theorem 3.5 we see that φ is the canonical morphism and X is irregular. $\square$

Corollary 3.7. *Let W be a smooth rational scroll of degree r and let $X \xrightarrow{\varphi} W$ be a Galois canonical cover with Galois group $\mathbf{Z}_4$ (i.e., a cover like the ones classified in Theorem 3.5 and Theorem 3.6). Then X is singular. Moreover,*

- 1) *if X is regular, then the mildest possible set of singularities on X consists of $4(r+1)$ singular points of type A_1 and*
- 2) *if X is irregular the singularities of X are exactly $4(r+4)$ points of type A_1 .*

Proof. The proof goes as the proof of Corollary 2.12, 3). In this occasion, $D_1 \cdot D_2 = 4(r+1)$ if X is regular (see Theorem 3.5, 1) and 2)) and $D_1 \cdot D_2 = 4(r+4)$ (see Theorem 3.6, 1) and 2)). X has the mildest possible set of singularities if X_1 is smooth and the branch locus of $X \xrightarrow{p_2} X_1$ has the mildest possible set of singularities. This happens if D_1 and D_2 are smooth and meet transversally. In this case the branch locus of p_2 has only singularities of type A_1 , and so does X . Now, if X is irregular Theorem 3.6 together with the fact that X is normal implies that D_1 is a union of distinct lines of one of the fibrations of $\mathbf{P}^1 \times \mathbf{P}^1$ and D_2 is a union of distinct lines of the other fibration, so D_1 and D_2 are smooth and meet transversally in any case. $\square$

Now we proceed to classify quadruple Galois canonical covers of smooth rational normal scrolls with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$. Having in account Proposition 2.6 we already know that they are the fiber product of two double covers. Thus to complete their description we will find out what the branch loci of the double covers are. We start with the case where X is regular:

Theorem 3.8. *Let $W = S(m-e, m)$ be a smooth rational normal scroll. If X is regular and $X \xrightarrow{\varphi} W$ is a Galois canonical cover with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$, then X is the fiber product over W of two double covers of $X_1 \xrightarrow{p_1} W$ and $X_2 \xrightarrow{p_2} W$ and φ is the natural map from the fiber product to W . Let the branch divisors D_2, D_1 of p_1, p_2 be linearly equivalent to $2a_2C_0 + 2b_2f$ and $2a_1C_0 + 2b_1f$ respectively. Then $0 \leq e \leq 2$, $m \geq e+1$, $a_1 = 1, a_2 = 2, b_1 = m+1$ and $b_2 = e+1$.*

Conversely, let $W = S(m, m-e)$ be such that $0 \leq e \leq 2$ and $m \geq e+1$ and let $X \xrightarrow{\varphi} W$ be the natural map to W from the fiber product over W of two flat double covers p_1 and p_2 with branch divisors as described above. Then X is regular and $X \xrightarrow{\varphi} W$ is a Galois canonical cover with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$.

Proof. Since X is regular, Proposition 1.6, 3) yields

$$\begin{aligned} \varphi_* \mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-C_0 - (m+1)f) \oplus \\ &\mathcal{O}_W(-2C_0 - (e+1)f) \oplus \mathcal{O}_W(-3C_0 - (m+e+2)f) . \end{aligned}$$

Then Proposition 2.6 tells us that X is the fiber product over W of two double covers of $X_1 \xrightarrow{p_1} W$ and $X_2 \xrightarrow{p_2} W$ with trace zero modules $L_1^* = \mathcal{O}_W(-2C_0 - (e+1)f)$ and $L_2^* = \mathcal{O}_W(-C_0 - (m+1)f)$ respectively, or equivalently, branched along divisors $D_2 \sim 4C_0 + 2(e+1)f$ and $D_1 \sim 2C_0 + 2(m+1)f$ respectively. Thus $a_1 = 1, a_2 = 2, b_1 = m+1$ and $b_2 = e+1$. Recall that W is isomorphic to the Hirzebruch surface $\mathbf{F}_e$. Since W is

smooth, $m \geq e + 1$, hence the only thing left to prove is $e \leq 2$. The covers p_1 and p_2 fit in the commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{p'_1} & X_2 \\ \downarrow p'_2 & & \downarrow p_2 \\ X_1 & \xrightarrow{p_1} & W \end{array}$$

where p'_1 and p'_2 are also double covers. Moreover the branch divisor of p'_1 is $p_2^* D_2$. Suppose that $e \geq 3$. Then, since D_2 is linearly equivalent to $4C_0 + (2e + 2)f$, D_2 has $2C_0$ as a fixed component. Then the branch divisor of p'_1 is nonreduced, so X is non-normal and we get a contradiction. Therefore $e = 0, 1$ or 2 .

To prove the converse assume now that $X \xrightarrow{\varphi} W$ is the natural map from the fiber product over a smooth scroll $W = S(m, m - e)$ of two double covers p_1 and p_2 of W , branched respectively along divisors D_2 linearly equivalent to $2a_2C_0 + 2b_2f$ and D_1 linearly equivalent to $2a_1C_0 + 2b_1f$. Assume in addition that $0 \leq e \leq 2$, $a_1 = 1, a_2 = 2, b_1 = m + 1$ and $b_2 = e + 1$. Then by Proposition 2.9 φ is a Galois cover with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$ and

$$\begin{aligned} \varphi_* \mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-a_1C_0 - b_1f) \oplus \\ &\mathcal{O}_W(-a_2C_0 - b_2f) \oplus \mathcal{O}_W(-(a_1 + a_2)C_0 - (b_1 + b_2)f) . \end{aligned}$$

A standard computation shows that none of the four direct summands of $p_* \mathcal{O}_X$ have intermediate cohomology, hence $H^1(\mathcal{O}_X) = 0$. On the other hand if $L_2 = \mathcal{O}_W(a_1C_0 + b_1f)$ and $L_1 = \mathcal{O}_W(a_2C_0 + b_2f)$, $L_1 \otimes L_2 = \mathcal{O}_W(3C_0 + (m + e + 2)f) = \omega_W^*(1)$. Then by Proposition 2.9 $\omega_X = \varphi^* \mathcal{O}_W(1)$ so X is a surface of general type whose canonical bundle is base-point-free. The only thing left to be shown is that $X \xrightarrow{\varphi} W$ is the canonical morphism of X . For that it is enough to see that $H^0(\mathcal{O}_W(1)) = H^0(\omega_X)$. But

$$\begin{aligned} H^0(\omega_X) &= H^0(\varphi^* \mathcal{O}_W(C_0 + mf)) = H^0(\mathcal{O}_W(C_0 + mf)) \oplus \\ &H^0(\mathcal{O}_W((1 - a_1)C_0 + (m - b_1)f)) \oplus H^0(\mathcal{O}_W((1 - a_2)C_0 + (m - b_2)f)) \oplus \\ &H^0(\mathcal{O}_W(1 - a_1 - a_2)C_0 + (m - b_1 - b_2)f)) . \end{aligned}$$

Now, because of the restrictions on a_1, a_2, b_1 and b_2 , $H^0(\mathcal{O}_W((1 - a_1)C_0 + (m - b_1)f))$, $H^0(\mathcal{O}_W((1 - a_2)C_0 + (m - b_2)f))$ and $H^0(\mathcal{O}_W(1 - a_1 - a_2)C_0 + (m - b_1 - b_2)f)$ vanish. $\square$

Now we go on to classify Galois quadruple covers with group $\mathbf{Z}_2 \times \mathbf{Z}_2$ when X is irregular:

Theorem 3.9. *Let W be a smooth rational normal scroll $S(m - e, m)$. If X is irregular and $X \xrightarrow{\varphi} W$ is a Galois canonical cover with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$, then X is the fiber product over W of two double covers of $X_1 \xrightarrow{p_1} W$ and $X_2 \xrightarrow{p_2} W$ and φ is the natural map from the fiber product to W . Let the branch divisors D_2, D_1 of p_1, p_2 be linearly equivalent to $2a_2C_0 + 2b_2f$ and $2a_1C_0 + 2b_1f$ respectively. Then $e = 0, m \geq 1$ and one of the following happens:*

- 1) $a_1 = 0, a_2 = 3, b_1 = m + 1, b_2 = 1$.

- 2) $a_1 = 0, a_2 = 3, b_1 = m + 2, b_2 = 0$.
 3) $a_1 = 1, a_2 = 2, b_1 = m + 2, b_2 = 0$.

In addition, in case 1), $q(X) = m$; in case 2), $q(X) = m + 3$; and in case 3), $q(X) = 1$.

Conversely, let $X \xrightarrow{\varphi} W$ be the natural map to $W = S(m, m)$ from the fiber product over W of two flat double covers p_1 and p_2 with branch divisors satisfying 1), 2) or 3) above. Then X is irregular and $X \xrightarrow{\varphi} W$ is a Galois canonical cover with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$.

Proof. From Proposition 1.3 and Proposition 2.6 it follows that X is the fiber product of two double covers branched along divisors $D_2 \sim 2(a_2C_0 + b_2f)$ and $D_1 \sim 2(a_1C_0 + b_1f)$ respectively, that

$$\begin{aligned} \varphi_*\mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-a_1C_0 - b_1f) \oplus \\ &\mathcal{O}_W(-a_2C_0 - b_2f) \oplus \mathcal{O}_W(-(a_1 + a_2)C_0 - (b_1 + b_2)f) \end{aligned}$$

and that $\omega_W(-1) = \mathcal{O}_W(-(a_1 + a_2)C_0 - (b_1 + b_2)f)$. Since $\omega_W = \mathcal{O}_W(-2C_0 - (e + 2)f)$, we obtain

$$\begin{aligned} a_1 + a_2 &= 3 \\ b_1 + b_2 &= m + e + 2 \end{aligned} \quad (3.9.1)$$

On the other hand since D_i is effective and linearly equivalent to $2(a_iC_0 + b_if)$, then $a_i, b_i \geq 0$. We set $a_1 = 0, 1$ (in which case, $a_2 = 3, 2$). Since φ is induced by the complete linear series $\varphi^*\mathcal{O}_W(C_0 + mf)$, then $H^0(\mathcal{O}_W((1 - a_1)C_0 + (m - b_1)f)) = 0$, hence $b_1 \geq m + 1$ and from (3.9.1), $b_2 \leq e + 1$. Since both b_1 and b_2 are nonnegative, $m + 1 \leq b_1 \leq m + e + 2$ and $0 \leq b_2 \leq e + 1$.

Now assume $a_1 = 0$. Then D_2 is linearly equivalent to $6C_0 + 2b_2f$. Assume also that $e \geq 1$. Then $2C_0$ is a fixed component of D_2 , therefore by the argument made in the proof of Theorem 3.8, X would be nonnormal, hence, if $a_1 = 0$, then $e = 0$. Then, we have two possibilities: first, $b_1 = m + 1$ and $b_2 = 1$, and second, $b_1 = m + 2$ and $b_2 = 0$. In the first case, $q(X) = m$. In the second case, $q(X) = m + 3$.

Now assume $a_1 = 1$. Then D_2 is linearly equivalent to $4C_0 + 2b_2f$, and as we argued in the proof of Theorem 3.8, if $e \geq 3$, X would be nonnormal, hence $0 \leq e \leq 2$. Moreover, if $b_2 < \frac{3e}{2}$, D_2 would have $2C_0$ as fixed component and X would be nonnormal, hence $b_2 \geq \frac{3e}{2}$. Recall also that $b_2 \leq e + 1$. Let us now assume that $b_2 = e + 1$. Then $b_1 = m + 1$ and in that case

$$\begin{aligned} \varphi_*\mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-C_0 - (m + 1)f) \oplus \\ &\mathcal{O}_W(-2C_0 - (e + 1)f) \oplus \mathcal{O}_W(-3C_0 - (m + e + 2)f) . \end{aligned}$$

But then $H^1(\mathcal{O}_W)$, $H^1(\mathcal{O}_W(-C_0 - (m + 1)f))$, $H^1(\mathcal{O}_W(-2C_0 - (e + 1)f))$ and $H^1(\mathcal{O}_W(-3C_0 - (m + e + 2)f))$ all vanish, hence X would be regular. Therefore $\frac{3e}{2} \leq b_2 \leq e$. This implies that $e = 0$ and $b_2 = 0$, in which case $b_1 = m + 2$. Then $q(X) = 1$. With this we prove that $W = S(m, m)$ and that the only possibilities for a_i s, b_i s are 1), 2) and 3).

To prove the converse, assume now that $X \xrightarrow{\varphi} W$ is the fiber product over W of two double covers p_1 and p_2 of W , branched respectively along divisors D_2 linearly equivalent to $2a_2C_0 + 2b_2f$ and D_1 linearly equivalent to $2a_1C_0 + 2b_1f$ satisfying one of the cases 1), 2) or 3). Then, by Proposition 2.9, φ is a Galois cover with Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$ and

$$\begin{aligned} \varphi_*\mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-a_1C_0 - b_1f) \oplus \\ &\mathcal{O}_W(-a_2C_0 - b_2f) \oplus \mathcal{O}_W(-(a_1 + a_2)C_0 - (b_1 + b_2)f) \end{aligned}$$

with a_1, a_2, b_1, b_2 satisfying 1), 2) or 3). Computing the cohomology of the four direct summands of $\varphi_*\mathcal{O}_X$ in each case shows that X is always irregular. We are going to see that $\omega_X = \varphi^*\mathcal{O}_W(1)$. This follows from Proposition 2.9 as in the proof of Theorem 3.8. Then in particular ω_X is base-point-free and X is a surface of general type. The only thing left to be shown is that $X \xrightarrow{\varphi} W$ is the canonical morphism of X . For that it is enough to see that $H^0(\mathcal{O}_W(1)) = H^0(\omega_X)$. This follows because $H^0(\omega_X) = H^0(\varphi_*\varphi^*\mathcal{O}_W(1))$, which is equal to $H^0(\mathcal{O}_W(1))$ by the same computation as in the proof of Theorem 3.8. $\square$

In the next corollary of Theorems 3.5, 3.6, 3.8 and 3.9 (see also Proposition 1.6, 3) if X is regular) we summarize the splitting of $\varphi_*\mathcal{O}_X$ if $X \xrightarrow{\varphi} W$ is a Galois canonical cover of degree 4 of a smooth rational normal scroll:

Corollary 3.10. *Let $X \xrightarrow{\varphi} W$ be a canonical Galois cover of degree 4, with Galois group G , of a smooth rational normal scroll isomorphic to $\mathbf{F}_e$ and embedded by $|C_0 + mf|$.*

1) *If X is regular, then $e = 0, 1, 2$ and*

$$\begin{aligned} \varphi_*\mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-C_0 - (m+1)f) \oplus \\ &\mathcal{O}_W(-2C_0 - (e+1)f) \oplus \mathcal{O}_W(-3C_0 - (m+e+2)f) . \end{aligned}$$

2) *If X is irregular, then $e = 0$ and*

$$\begin{aligned} 2.1) \quad \varphi_*\mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-C_0 - (m+2)f) \oplus \mathcal{O}_W(-2C_0) \oplus \\ &\mathcal{O}_W(-3C_0 - (m+2)f) ; \text{ or} \end{aligned}$$

$$\begin{aligned} 2.2) \quad \varphi_*\mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-(m+1)f) \oplus \mathcal{O}_W(-3C_0 - f) \oplus \\ &\mathcal{O}_W(-3C_0 - (m+2)f) , \text{ with } m = 1 \text{ if } G = \mathbf{Z}_4; \text{ or} \end{aligned}$$

$$\begin{aligned} 2.3) \quad \varphi_*\mathcal{O}_X &= \mathcal{O}_W \oplus \mathcal{O}_W(-(m+2)f) \oplus \mathcal{O}_W(-3C_0) \oplus \mathcal{O}_W(-3C_0 - (m+2)f), \\ &\text{and } G = \mathbf{Z}_2 \times \mathbf{Z}_2. \end{aligned}$$

We end the section showing the existence of canonical covers like the ones classified in Theorems 3.5, 3.6, 3.8 and 3.9:

Proposition 3.11. *There exist families of quadruple Galois canonical covers as in Theorem 3.5 and Theorem 3.6 which have singularities as mild as possible (see Corollary 3.7).*

Proof. According to the converse part in Theorem 3.5 and Theorem 3.6 we just have to construct a composition of double covers $X \xrightarrow{p_2} X_1$ and $X_1 \xrightarrow{p_1} W$ branched along suitable divisors. Let D_1 and D_2 be as in Theorem 3.5 and Theorem 3.6. By the arguments of

Corollary 2.12 and Proposition 2.13, in order for X to have singularities as mild as possible and, in any case, only A_1 singularities, it suffices to choose D_1 and D_2 smooth and meeting transversally. We see that such choice is indeed possible. For a cover as in Theorem 3.5, 1), $D_1 \sim (2m - e + 1)f$ and $D_2 \sim 4C_0 + (2e + 2)f$. Then we choose D_1 as the union of $2m - e + 1$ different fibers. The divisor $4C_0 + (2e + 2)f$ is base-point-free if $e = 0, 1$ and if $e = 2$ is $(3C_0 + 6f) + C_0$, with $3C_0 + 6f$ base-point-free and $(3C_0 + 6f) \cdot C_0 = 0$. Thus by Bertini D_1 and D_2 can be chosen smooth and intersecting transversally. For a cover like Theorem 3.5, 2), both $D_1 \sim 3C_0$ and $D_2 \sim 2C_0 + 4f$ are base-point-free (recall that in this case W is isomorphic to $\mathbf{F}_0$), so we can apply Bertini as before. Finally, in Theorem 3.6 D_1 and D_2 are the union of distinct smooth lines belonging to the two fibrations of the ruled surface $\mathbf{F}_0$. Note that, if X is regular, one can construct X with worse singularities allowing $D_1 + D_2$ to have worse singularities. $\square$

Proposition 3.12. *There exist families of canonical Galois quadruple covers $X \xrightarrow{\varphi} W$ as in Theorem 3.8 and Theorem 3.9, 1), 2) and 3) with X smooth.*

Proof. Families satisfying Theorem 3.8 for W isomorphic to $\mathbf{F}_0$ and $\mathbf{F}_1$ have been constructed in [GP2], Examples 3.4 and 3.5 (see also [Pe] for an example of a bidouble cover of $\mathbf{P}^1 \times \mathbf{P}^1$). To construct the remaining examples we argue as in Proposition 2.13. By the converse part of Theorem 3.8 and Theorem 3.9 we just need to construct the fiber product of two double covers $X_1 \xrightarrow{p_1} W$ and $X_2 \xrightarrow{p_2} W$ branched along suitable divisors D_2 and D_1 satisfying the conditions in the statement of Theorem 3.8 and Theorem 3.9. Precisely if we choose D_1 and D_2 smooth and meeting transversally, X will be smooth. This can be achieved using Bertini once we study how the divisors D_1 and D_2 are in each case. Indeed, if we are in the situation of Theorem 3.8 when W is isomorphic to $\mathbf{F}_2$, then $D_1 \sim 2C_0 + 2(m + 1)f$ is very ample and D_2 , since it is linearly equivalent to $4C_0 + 6f$ is of the form $C_0 + D'_2$, with $D'_2 \cdot C_0 = 0$ and D'_2 base-point-free.

In case Theorem 3.9, 1), 2) and 3) recall that W is isomorphic to $\mathbf{F}_0$. In case Theorem 3.9, 1) $D_1 \sim 2(m + 1)f$, hence it can be chosen as the union of $2(m + 1)f$ distinct lines in one of the two fibrations of $\mathbf{F}_0$ and D_2 is linearly equivalent to $6C_0 + 2f$, which is very ample.

Finally in case Theorem 3.9, 3), $D_1 \sim 2C_0 + 2(m + 2)f$ is very ample and $D_2 \sim 4C_0$ can be chosen as the union of 4 distinct lines of one of the fibrations of $\mathbf{F}_0$. Note that one can construct X with worse singularities if $D_1 + D_2$ is allowed to have worse singularities. $\square$

REFERENCES

- [Be] A. Beauville, *L'application canonique pour les surfaces de type general*, Inventiones Math. **55** (1979), 121-140.
- [BS] M.C. Beltrametti and T. Szemberg, *On higher order embeddings of Calabi-Yau threefolds*, Arch. Math. (Basel) **74** (2000), 221-225.
- [C] F. Catanese, *On the moduli spaces of surfaces of general type*. J. Differential Geom. **19** (1984), no. 2, 483-515.
- [GP1] F.J. Gallego and B.P. Purnaprajna, *Very ampleness and higher syzygies for Calabi-Yau threefolds*, Math. Ann. **312** (1998), 133-149.

- [GP2] F.J. Gallego and B.P. Purnaprajna, *On the canonical ring of covers of surfaces of minimal degree*, Trans. Amer. Math. Soc. **355** (2003), 2715-2732.
- [GP3] F.J. Gallego and B.P. Purnaprajna, *Classification of quadruple canonical covers: Galois case*, to appear in C. R. Math. Acad. Sci. Soc. R. Can.
- [GP4] F.J. Gallego and B.P. Purnaprajna, *Classification of quadruple Galois canonical covers, II*, preprint.
- [GP5] F.J. Gallego and B.P. Purnaprajna, *Ring generation for cyclic and quadruple canonical covers*, in preparation.
- [Ga] X. Gang, *Algebraic surfaces with high canonical degree*, Math. Ann, **274** (1986), 473-483.
- [Gr] M.L. Green, *The canonical ring of a variety of general type*, Duke Math. J. **49** (1982), 1087-1113.
- [HM] D. Hahn and R. Miranda, *Quadruple covers of algebraic varieties*, J. Algebraic Geom. **8** (1999), 1-30.
- [Ho1] E. Horikawa, *Algebraic surfaces of general type with small c_1^2 , I*, Ann. of Math. (2) **104** (1976), 357-387.
- [Ho2] E. Horikawa, *Algebraic surfaces of general type with small c_1^2 , II*, Invent. Math. **37** (1976), 121-155.
- [Ho3] E. Horikawa, *Algebraic surfaces of general type with small c_1^2 , III*, Invent. Math. **47** (1978), 209-248.
- [Ho4] E. Horikawa, *Algebraic surfaces of general type with small c_1^2 , IV*, Invent. Math. **50** (1978/79), 103-128.
- [K] K. Konno, *Algebraic surfaces of general type with $c_1^2 = 3p_g - 6$* , Math. Ann. **290** (1991), 77-107.
- [MP] M. Mendes Lopes and R. Pardini, *Triple canonical surfaces of minimal degree*, International J. Math. **11** (2000), 553-578.
- [OP] K. Oguiso and T. Peternell, *On polarized canonical Calabi-Yau threefolds*, Math. Ann. **301** (1995), 237-248.
- [Pa] R. Pardini, *Abelian covers of algebraic varieties*, J. Reine Angew. Math. **417** (1991), 191-213.
- [Pe] U. Persson, *Double coverings and surfaces of general type*, Algebraic Geometry. (Lect. Notes Math., vol 687. 168-195) **687** (1978).
- [Pu] B.P. Purnaprajna, *Geometry of canonical covers with applications to Calabi-Yau threefolds*, in Vector Bundles and Representation Theory, Eds. D. Cutkosky et al., Contemporary Mathematics Series AMS **322** (2003), 107-124.
- [T] S. Tan, *Surfaces whose canonical maps are of odd degrees*, Math. Annalen, **292** (1992), 13-29.

FRANCISCO JAVIER GALLEGO: DPTO. DE ÁLGEBRA, FACULTAD DE MATEMÁTICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, 28040 MADRID, SPAIN
E-mail address: gallego@mat.ucm.es

B.P.PURNAPRAJNA: 405 SNOW HALL, DEPT. OF MATHEMATICS, UNIVERSITY OF KANSAS, LAWRENCE, KANSAS 66045-2142
E-mail address: purna@math.ukans.edu